



Effect of *Azospirillum zae* and Seed Priming with Zinc, Manganese and Auxin on Growth and Yield Parameters of Wheat, under Dryland Farming

Nasrin Karimi¹ · Ebrahim Mohammadi Goltapeh² · Jahanshir Amini³ · Samina Mehnaz⁴ · Mohammad Javad Zarea¹

Received: 5 June 2019 / Accepted: 20 July 2020

© NAAS (National Academy of Agricultural Sciences) 2020

Abstract In this study, field experiments with wheat (*Triticum aestivum* L.) were conducted consecutively for 2 years. The main objective was to determine (1) effect of seed priming with auxin (IAA), zinc (Zn) or manganese (Mn) on growth and yield parameters of wheat under dry land farming; (2) influence of on-farm priming with *Azospirillum* under field conditions. In this study, the response of dryland wheat yield to soaking seeds in water, 2 ppm IAA, 0.2% Mn solution and 0.2% aqueous solution of Zn with or without *Azospirillum zae* inoculation was investigated under field condition. Total grain yield was significantly enhanced for water primed seeds i.e., 1273 kg ha⁻¹ (6.8%) or seeds primed with IAA, as compared to 1191 kg ha⁻¹ in plants from non-primed seeds. Difference in grain yield between priming with water and 2 ppm IAA was not significant when compared with 1372 kg ha⁻¹ (15%) for seeds primed with 0.2% Mn solution and 1440 kg ha⁻¹ (20%) for seeds primed with 0.2% Zn solution. Grain yield was significantly increased for plants inoculated with *Azospirillum*, 1411 kg ha⁻¹ (8%) as compared to 1191 t ha⁻¹ for non-*Azospirillum*-inoculated seeds. Yield parameters, i.e., tiller no. per m⁻², grain number per spike, and 1000-grain weight, were all significantly affected by priming treatments and *Azospirillum* inoculation. Grains with the high nitrogen and phosphorous content were recorded for the treatment where seeds were primed with Zn and inoculated with *Azospirillum*. In the present study, inoculation of wheat seeds followed by priming is a significant approach to enhance grain yield and grain nutrients (N and P) content.

Keywords Seed priming · Micronutrients · Auxin · *Azospirillum*

Introduction

On-farm seed priming has potential to improve grain yield. This technique has been adopted by resource-poor farmers for a verity of crops. Plant-growth-promoting bacteria, such as *Azospirillum*, have also been frequently reported that

have potential to increase crop yield both under non-stress and stress conditions.

Various macro- and micronutrients are necessary for the proper plant production. In most agricultural lands, farmers tend to use macronutrients such as N, P and K to gain the higher plant yields, while micronutrients such as Fe, Zn, and Mn have been reported to improve crop growth, grain nutrient enrichment and plant establishment under farm condition [23]; however, applying these micronutrients is not common a practice by farmers in arable land as compared to macronutrients.

Micronutrients play vital role in plants as they are involved in several biological processes. Zn is involved in several biological functions and metabolic reactions [79]. Plants require a low amount of Fe and Zn for their physiological and metabolic processes. Their excess or deficit of these micronutrients induces negative consequences in

✉ Mohammad Javad Zarea
mjzarea@yahoo.com; mj.zarea@ilam.ac.ir

¹ Faculty of Agriculture, Ilam University, Ilam, Iran

² Department of Plant Pathology, Agriculture Faculty, Tarbiat Modares University, P.O. Box 14115-336, Tehran, Iran

³ Department of Plant Protection, Faculty of Agriculture, University of Kurdistan, Sanandaj, Iran

⁴ Department of Biological Sciences, Forman Christian College (A Chartered University), Lahore, Pakistan

leaves, root system, plant weight, overall biomass, photosynthesis, can damage the DNA, and directly affect the cell cycle and chromosomes [45], [56], [63], [46], [68].

Mn is involved in photosynthesis and nitrogen metabolism [82], [76]. More than 300 plant enzymes and some vital proteins have Zn in their structure [88], [43]. Plant requirement for micronutrients is low, and excess of these element results in damages to plant such as excessive concentration of Zn in plant causes damage to the DNA, cell cycle and chromosomes [45], [29], [46].

Wheat, as the most important crop, provides nutrition to most of the world population [24] and plays important role in human diet. Deficit of Zn and Mn in human diet is responsible for some diseases [12,14]. Transgenesis and classic breeding methods have been used to improve the micronutrient content of modern wheat cultivars [68]; however, other methods, such as foliar spraying, seed priming and soil application, have been shown to be cost-effective in this regard [26], [9], [23], [77]. Priming seeds among the above-mentioned methods has been reported to be the most effective in improving crop yield as well as the most cost-effective method in increasing micronutrient levels in plants such as wheat [78], [23]. In seed priming, seeds are allowed to be partially hydrated (hydro-priming) to trigger metabolic events without actual germination occurring and then re-dried to attain their near original weight [8]. Primed seeds usually have better germination and establishment, weight and height, root length and grain yield [32], [23], [68].

Improvement in grain yield of wheat followed by priming with Zn has been reported by Padole [61], [75], and Harris et al. [33]. Zn deficiency restricts the growth and yield of wheat [33]. Zn deficiency have been reported to be common in developed and developing countries [92] and, for instant, mentioned as the third most nutrition problem in Pakistan after N and P deficiency [67]. Seeds with higher level of micronutrients have promising results. It had been reported earlier that elevating Zn content in wheat seed improved seedling vigor and enhanced yield [93], [33]. Incidences of micronutrient deficiency in crops such as Mn and Zn seem to have enhanced worldwide in recent years [17]. Garvin et al. [25] reported that Zn contents in wheat grain are inversely correlated with yield. Previous studies showed that use of seeds with elevated micronutrient content resulted in enhanced grain yield and improved seedling vigor [69, 70], [93]. Reis et al. [68] investigated the effect of seed priming with different concentrations of Zn (1–8 mg L⁻¹) on germination and yield of wheat. Result of this experiment showed that, however, Zn dosages higher than 8 mg L⁻¹ had negative effects on germination and grain yield, but seed priming with lower dosages of Zn was non-cytotoxic and improved rate of germination and grain yield.

Among plant hormones, auxins were the first one to be discovered. Indole-3-acetic acid (IAA) is the most prevalent plant hormone of auxin class. IAA is critical for the regulation of the various aspects of plant growth and development [86] as it plays vital role in cell division and differentiation [73]. Lecube et al. [44] reported the role of IAA in protecting plants against oxidative stress in soybean. Environmental factors affect adversely such as drought effect on synthesis of some plant hormones, which causes limitation on physiological processes of the plant [16]. Exogenous application of these hormones at lower concentration regulates differentiation, growth, and development of the plant [55]. The application of these hormones causes the physiological processes to occur at their normal rate [28]. Auxin induces root initiation, and development which can be important under drought stress condition. Improved root system can assist plant in exploring more moisture from depth of soil and enhances absorption of the nutrients as well. Priming seeds with plant growth regulators have been suggested to enhance seed germination, seedling growth, and the yield of plant [22], [21], [91]. Priming seeds of corn [62], wheat [87], rice [91], beet [21], sunflower [36], safflower [3], and perennial grass [47] with plant growth regulators have been reported to result in enhancing yield.

Nitrogen fixing bacteria are frequent colonizer of cereal crops and grasses belonging to the genera *Azospirillum*, *Acetobacter*, *Azoarcus*, *Enterobacter*, and *Herbaspirillum* [50]. The genus *Azospirillum* was first reported by Beijerinck [6] as *Spirillum* and later by the group of Dr. Johanna Döbereiner in Brazil, in the 1970s [85] reclassified as *Azospirillum*. It is a diazotrophic bacteria associated with a wide range of plants. A variety of *Azospirillum* species and other PGPR have been reported to have ability to colonize the aboveground parts and root systems of various plants such as maize [51], [52], wheat [42], rice [41], sugarcane, and other plants [81]. Growth-promoting effect of *Azospirillum* on yield of cereals such as wheat under field conditions has been elucidated [72], [35], [96], [38]. In cereals, *Azospirillum* resulted in increased plant biomass, nutrient uptake and root length and volume [72]. Phytohormone production is one of the mechanisms proposed for *Azospirillum* by which it improves plant growth [57], [5]. Aside from this benefit, *Azospirillum* directly contributes to make N available to the plants [18].

The current study aimed to assess the growth and yield response of wheat to different seed priming treatments, i.e., Zn, Mn, IAA and distilled water, with or without *Azospirillum zeae* under dryland farming condition. A preliminary experiment was also conducted to determine the optimum concentration of priming solutions of Mn and Zn, with and without inoculation of *Azospirillum*. *Azospirillum* used in this experiment was a native strain,

adapted to the region where this experiment was conducted. Numerous reports have been shown that inoculation with *Azospirillum* resulted in improvement of plant growth and yield of wheat [96]. PGPR can promote plant growth of the host plant through different mechanisms such as biological nitrogen fixation, production of indolic compounds, siderophore production, ACC deaminase activity, phosphate solubilization and plant anatomy modulation [57], [5], [19], [94], [95], [35], [97], [98]. Appreciate agronomical and technical management can affect PGPR performances [96]. In this study, seed priming as an agronomical management was used to investigate whether it can enhance promoting effect of *Azospirillum* on wheat under dryland farming.

Materials and Methods

Bacterial Strains

Indigenous bacteria used in this study were previously isolated by Dr. MJ. Zarea. It was isolated from the inside of roots of wild wheat, naturally growing in semiarid mountainous areas of west Iran. Preliminary studies conducted in our laboratory indicated that this strain was capable of IAA synthesis and nitrogen fixation. This strain, *Azospirillum zeae* MK433605, was able to produce indolic compounds at the rates of $37.46 \pm 0.84 \mu\text{g ml}^{-1}$. The ability of bacterial isolate to solubilize phosphate was $1.89 \pm 0.14 \text{ mg L}^{-1}$. Identification of the species of the bacterial isolate based on the molecular approaches was done by Dr S. Mehnaz. Based on the 16S rRNA gene sequence analyses, bacterial strain was identified as *Azospirillum zeae*, (> 98% similarity with *A. zeae*). The 16S rRNA gene sequence was deposited in the NCBI databank, getting the accession numbers MK433605. For seed priming with bacterial strain, culture of bacterial strain was prepared in 1 L of NFB [38] using 1 mL of the overnight-grown bacterial culture as initial inoculum.

Preliminary Experiments

The optimization of concentration of Zn and Mn priming solution with and without inoculation of *Azospirillum* was done in pot culture, with the dosage information available in Johnson et al. [37] and [33] for Zn and [39] for Mn. Wheat seeds (*Triticum aestivum* L.) cultivar pishgam were first surface sterilized using 98% ethanol for 30 min, 2% sodium hypochlorite for 3 min and, then, washed thoroughly with distilled water. Seeds were primed for 6 h with aqueous solutions of ZnSO_4 or MnSO_4 at four concentrations, each of Zn and Mn, 0.1%, 0.2%, 0.3% and 0.4%.

Non-primed seeds and seeds primed with water served as control treatments.

All seeds from priming treatments and non-primed seeds were divided into two groups and each group of seeds received 1 mL of 2.0×10^7 CFU/g seed of *Azospirillum zeae*, inoculum or heat-killed *Azospirillum* inoculum (control), and, then, air-dried in the dark for 2 h at 25 °C.. Fifteen air-dried seeds from each priming treatment inoculated with or without *Azospirillum* were sown at the depth of 3 cm in plastic pot filled with a field soil and then each pot were thinned to maintain 10 plants per pot. The pots were placed in a greenhouse with natural light. At 30 day after sowing, seedlings of each pot were harvested and then top dry mass was measured. The experiment was a 6×2 factorial, arranged in a randomized block design with three replications of each treatment. Effects of seed priming and *Azospirillum* inoculation, and relevant interaction were analyzed, using analysis of variance, by SAS Software version 9.1.

Field Study

A field experiment was conducted for 2 years (2016–17) at agriculture land in Dehgolan County, Kurdistan Province ($35^{\circ}17'N$, $47^{\circ}22'E$, West Iran), on clay soil. The preceding crop was wheat in the previous year. Figure 1 shows the sum of rainfalls during the experiments. The experiment was a 5×2 factorial, arranged in a randomized complete block design with 3 replicates. Size of each plot was 3 m^2 , with 6 rows, each one 15 cm apart. Adjacent plots were separated by 1 m. No herbicides were applied before planting. Planting rate was 133 seeds m^{-2} . Chemical properties of the soil are presented in Table 1. The soil was slightly alkaline with low amount of organic matter. Seed beds were prepared during autumn. Sowing date was dependent on the time of first sufficient rainfall. A rate of 100 kg ha^{-1} nitrogen (urea) and phosphorus (50 kg ha^{-1}) in the form of triple superphosphate was applied on a day before sowing. Further 50 kg N ha^{-1} was applied as top dressing. Because of low density of weeds in plots, no foliar herbicide was applied and weeds were controlled through hand-weeding. Ten treatments included (1) Mn priming + bacterial inoculums (BI), (2) Zn priming + BI, (3) IAA priming + BI, (4) water priming + BI, (5) Mn priming, (6) Zn priming, (7) IAA priming, (8) water priming, (9) no priming, and (10) no inoculum, seeds as control were planted into plots, each with three replicates.

Seed Priming and *Azospirillum* Inoculation

Before starting the experiment, wheat seeds (*Triticum aestivum* L.) cv. pishgam were surface-sterilized using 98% ethanol for 30 min, 2% sodium hypochlorite for 3 min and




Fig. 1 Sum of rainfalls at 1-month intervals during the experiment

Table 1 Soil chemical properties in the first year of the establishment of the experiments

Soil pH	7.4
P, mg kg ⁻¹	12.29
K, mg kg ⁻¹	132
Zn, mg kg ⁻¹	0.76
Mg, mg kg ⁻¹	92
Mn, ppm	0.58
EC, (dS m ⁻¹)	1.82
Organic matter	1.67

then washed thoroughly with distilled water. For the hormo-priming, sterile seeds were soaked for 6 h, in 2 ppm solution of IAA and air-dried under shade, prior sowing. IAA stock was prepared by adding 0.002 g IAA to a 10 mL volumetric flask and dissolved by 95% ethanol, after which sterile distilled water was added to make a total solution of 2 ppm IAA. For nutrient and hydro-priming, surface sterile seeds were soaked in distilled water, Zn (0.2%) or Mn (0.2%) solution for 6 h and then air-dried under shade to original weight to facilitate clump-free sowing. ZnSO₄ and MnSO₄ were used to prepare the priming solution. Zn and Mn concentrations used in field experiments were based on preliminary experiments.

Primed seeds were then air-dried and inoculated with *Azospirillum*. Half of each primed seeds were inoculated with bacterial inoculum, and the rest received heat-killed inoculum. Non-primed, non-inoculated seeds were also planted as control treatment. *Azospirillum zeae* was grown in NFb medium overnight. In inoculated seeds, 1 g of wheat seeds was inoculated with 1 mL suspension of washed overnight bacterial culture at a concentration of 2.0×10^7 CFU mL⁻¹.

Seedling Growth and Seed Emergence

Field emergence of seeds (%) was evaluated 2 weeks after sowing. Plant samples were taken from each plot at 30th and 60th day after planting, for measurements of dry weight. Samples consisted of five randomly selected plants from each plot (fifteen plants per treatment). Plant samples were dried at 75 °C and then weighed.

Yield Component Traits

At maturity, yield components were determined from five randomly selected plants. Yield components included number of tillers, number of spikelet per spike, number of grains per spike and 1000 grain weight (g). Grain yield was recorded from 1 m² of the plot center. 1000 grain weight was measured from five randomly selected spikes. Total grain yield was presented as kg h⁻¹ using the following formulae (a).

$$\text{Grain yield (kg/ha)} = \left(\frac{\text{Grain yield}}{\text{Sampled area}} \right) \times 10,000 \text{ m}^2$$

Nitrogen, Phosphorus and Protein Content in Grains

The method described by Van Schouwenberg and Walinge [89] was followed to estimate the nitrogen content in grains. For the estimation of phosphorus contents, ammonium vanadate-molybdate method described by Chapman and Pratt [11] was followed. Flag leaf soluble proteins were determined according to the method described by Bradford [7].

Statistical Analysis

The data recorded on field emergence, seedling fresh weight, yield components, grain yield and P and N content of grain for 2 years were pooled to get a mean value. Effects of seed priming and *Azospirillum* inoculation, and relevant interaction on all data recorded were analyzed, using the PROC GLM procedure of SAS, by SAS software version 9.1. Differences among treatment means were identified using Fisher's protected LSD_{0.05} analysis.

Results

Preliminary Experiment

In preliminary experiment, 30 days after sowing, plants primed with 0.2% or 0.3% Zn showed significantly high dry weight than plants from non-primed seeds, primed with 0.1% Zn or 0.4% Zn (Fig. 2). Priming with 0.4% Zn reduced seedlings dry weight (Fig. 2). Dry weight of seedlings, at 30 days after sowing, in all plants from non-primed seeds and Zn-primed seeds became heavier due to *Azospirillum* inoculation. *Azospirillum* inoculation increased dry weight of 30-d old seedlings in non-primed plants and Zn-primed plants by 15.48% and by 26.47%, respectively. Treatment seeds with 0.2% Zn + *Azospirillum* inoculation significantly increased weight of plants as compared to other treatments (Fig. 2).

Priming seeds with 0.1%, 0.2%, 0.3% or 0.4% Mn increased dry weight of 30-day-old seedlings compared to non-primed seeds and seeds primed with water (Fig. 2). Using 0.4% Mn decreased dry weight of seedlings

compared to 0.2% or 0.1% Mn. Accompanied Mn priming with *Azospirillum* inoculation resulted in an increase in dry weight of 30 days old seedlings (Fig. 2). Plants from seeds primed with 0.2% or with 0.3% Mn and inoculated with *Azospirillum* had higher dry weight than those from non-primed seeds, primed with water and primed with 0.1%, or with 0.4% Mn (Fig. 2).

Field Experiments

Seed priming and *Azospirillum* inoculation significantly influenced ($P < 0.05$) all parameters recorded, including early fresh seedling weight, grain yield and components, and nutrient (N and P) uptakes (except for field emergence percent). In this study, field emergence percent was not significantly affected by all treatments of seed priming with or without *Azospirillum* inoculation in comparison with not primed, not inoculated seeds. Results of combined analysis of the 2-year field experiments indicated that priming seeds with water, IAA (2 ppm), MnSO₄ (0.2%) or with ZnSO₄ (0.2%) did not reduce yield, yield components and N and P content in grain.

Early Seedling Weight

In general, fresh weight of wheat seedlings was improved with priming treatments (Zn, Mn, IAA and water) or inoculated with *Azospirillum*, but *Azospirillum* inoculation combined with seed priming significantly elevated the plant biomass (fresh weight) at 30 and 60 days after sowing. Figure 3 shows the effect of seed priming and *Azospirillum* inoculation on fresh seedling weight at 30 and 60 days after sowing. Plants from seeds primed with Zn,




Fig. 2 The effect of priming wheat seeds for 6 h with various concentration rates (%) of aqueous solutions of Zn and Mn inoculated or not inoculated with *Azospirillum* on 30-day-old wheat seedlings

dry mass. Values with the same letters are not significantly different at $P < 0.05$. Values are mean \pm standard error (SE) ($n = 3$)




Fig. 3 The effect of priming seeds for 6 h in water, IAA (2 ppm), or with aqueous solutions of Zn (0.2%) or Mn (0.2%) on seedling fresh weight at 30 and 60 days after sowing (DAS). Values with the same

letters are not significantly different at $P < 0.05$. Values are mean \pm standard error (SE) ($n = 3$)

Mn, IAA or water were significantly heavier at 30 and 60 days after sowing than that of from non-primed seeds. Difference in effect of priming and *Azospirillum* inoculation was more visible at 60 days after sowing (Fig. 3). At 60 days after sowing, plants from seeds primed with 0.2% Zn and inoculated with *Azospirillum* were significantly heavier than those from non-primed seeds, Zn priming alone and from other seed priming with or without *Azospirillum* inoculation (Fig. 3).

Grain Yield

The interactive effect of seed priming and *A. zae* inoculation on wheat grain yield is presented in Fig. 4. The main effect of priming and *Azospirillum* inoculation and priming \times inoculation was significant for grain yield. In control plants, the grain yield was 1191 kg ha⁻¹ and enhanced to 1411 kg ha⁻¹ (18%) when inoculated with *Azospirillum*. Priming seed with Zn, Mg, IAA and water alone and in combination with *A. zae* inoculation significantly ($p < 0.05$) enhanced the grain yield of wheat compared to values for non-primed seeds (Fig. 4). Grain yield in plants primed for 6 h in 0.2% Zn or 0.2% Mn solution was significantly higher than that in plants primed with IAA and water (Fig. 4). Grain yield in primed seeds were increased with *Azospirillum* inoculation. Plants from Zn priming and *Azospirillum* inoculation produced the highest grain yield (Fig. 4). *Azospirillum* enhanced grain yield of plants (ha⁻¹)

from seeds primed in Zn, Mn, IAA and water by about 49%, 35%, 33% and 24%, respectively.

In this study, the benefits of seeds priming simply with water from those produced by added IAA, Mn and Zn were separated. Figure 5 shows the percent increase in grain yield of plants primed with IAA, Mn and Zn alone in combined or without *Azospirillum* inoculation over grain yield of plants from seeds primed with water. Grain yield was significantly increased by priming with water alone (1.06%) over the non-primed treatment. Grain yield was significantly increased further by adding small amount of Zn or Mn in water priming (Fig. 5). However, priming with 2 ppm IAA was not significantly superior to water priming (Fig. 5). Percent increase in grain yield was significantly higher when *Azospirillum* inoculation was accompanied with priming agent, IAA, Mn and Zn (Fig. 5). Supplying 0.2% Zn through water priming and then inoculation with *Azospirillum* gained an additional increase in grain yield of 35% as compared to water priming alone (Fig. 5).

Yield Components

Although the grain number per spike was not significantly affected by the seed priming treatments, it was significantly enhanced by *Azospirillum* inoculation (Table 2). Plants from seeds inoculated with *Azospirillum* produced, on average, 5.29% higher grain number per spike as compared to non-inoculation treatment. *Azospirillum* inoculation




Fig. 4 The effect of priming seeds for 6 h in water, IAA (2 ppm), or with aqueous solutions of Zn (0.2%) or Mn (0.2%) with or without *Azospirillum* inoculation on grain yield. Values with the same letters

are not significantly different at $P < 0.05$. Values are mean \pm standard error (SE) ($n = 3$)




Fig. 5 Separated benefits of seeds priming simply with water from those produced by added IAA, Mn and Zn coupled with or without *Azospirillum* inoculation. Values with the same letters are not significantly different at $P < 0.05$. Values are mean \pm standard error (SE) ($n = 3$)

resulted in improving effect on plants in terms of grain no. per spike. When seed priming combined with *Azospirillum* inoculation, enhancing in ration of grain no. per spike was more pronounced (Table 2). *Azospirillum* increased grain number per spike of plants from Zn, Mn, IAA and water priming treatments from 20.28 to 21.89 (7.93%), 20.11 to 22.64 (12.59%), 20.61 to 24.58 (19.27%) and 21.3 to 26.43 (24.09%), respectively. The highest grain number per spike was recorded for plants primed with 0.2% Zn solution for 6 h and inoculated with *Azospirillum* (Table 2).

Although 1000-grain weight was not significantly affected by the seed priming treatments, it was significantly enhanced by *Azospirillum* inoculation (Table 2). Plants from *Azospirillum*-inoculated seeds produced, on average, heavier 1000 grain weight (9%) than non-inoculated plants. There was a significant interaction between seed priming and *Azospirillum* inoculation on grain weight (Table 2). The greatest 1000 grain weight was obtained for plants, primed with Zn solution and inoculated with *Azospirillum* (Table 2).

Table 2 Comparison of seed priming with 0.2% Zn, 0.2% Mn and 2 ppm IAA with and without *Azospirillum* inoculation on yield components in wheat under dryland farming

Treatments		1000-seed weight (g)	Grain no spike ⁻¹	Spike no m ⁻²
With <i>Azospirillum</i>	Primed 0.2% Zn	32.72(± 1.32)a	26.43(± 1.26)a	250(± 5.3)a
	Primed 0.2% Mn	30.91(± 1.12)b	24.58(± 0.63)b	245(± 3.7)ab
	Primed 2 ppm IAA	30.84(± 0.89)b	22.64(± 0.5)cd	243(± 4.1)bc
	Primed H ₂ O	31.32(± 0.98)ab	21.89(± 0.33)d	231(± 5.6)d
	Not primed	29.46(± 1.02)bc	20.73(± 0.56)ef	230(± 4.5)d
Without <i>Azospirillum</i>	Primed 0.2% Zn	28.82(± 1.12)c	21.30(± 0.14)de	234(± 4.7)cd
	Primed 0.2% Mn	28.20(± 1.21)c	20.61(± 0.26)ef	236(± 3.8)c
	Primed 2 ppm IAA	27.36(± 1.54)c	20.11(± 0.24)ef	231(± 5.1)d
	Primed H ₂ O	27.65(± 1.32)c	20.28(± 0.8)ef	223(± 4.6)ef
	Not primed	27.01 ± (0.56)c	19.69(± 0.86)f	221(± 3.7)f

LSD_{0.05} for the *Azospirillum* inoculation × seed priming interaction

Values in parentheses are the standard deviation

Data (mean ± SE) are mean value of 2 years

Values with the same letters are not significantly different at $P < 0.05$

Values are mean ± standard error (SE) ($n = 3$)

Another yield component, spike number per m², was significantly affected by *Azospirillum* inoculation, but not by priming treatment. In control plants (non-priming) and in plants primed with water (hydro-priming), increase in number of spike per m² was the same and increased, on average, by 4.5% due to *Azospirillum* inoculation. The highest number of spike number per m² was recorded in plants inoculated with *Azospirillum* and primed with Zn.

Nitrogen and Phosphorus Content

Figure 6 shows a comparison of the grain N content of wheat among the priming treatments with and without *Azospirillum* inoculation. Grain N content in non-priming plants was 0.76% and in response to *Azospirillum* inoculation increased to 0.91%. Priming seeds with Zn and Mn had a positive effect on grain N content. Grain N content in plants from seeds primed in Zn and Mn solution increased by 20% and 15% compared with those of plants from non-priming seeds, IAA priming and hydro-priming. There was no significant difference between seed priming with water and IAA with control (not primed seeds) on N content of grain. Priming seeds in 2% Zn solution increased grain N content compared with all other priming treatments (Fig. 6). There was a significant interaction between seed priming and *Azospirillum* on grain N content. Grains with the highest N content obtained from plants underwent Zn priming and *Azospirillum* inoculation (Fig. 6).

Grain P content followed the same trend as observed for grain N content (Fig. 6). P content of grain was increased in non-primed treatment by *Azospirillum* inoculation,

5.93%. In this study, Zn priming among priming treatments was the most important influence on grain P content. Plants from seeds primed in 0.2% Zn solution for 6 h produced the grains with higher P content compared with those of plants from all the other priming treatment (Fig. 6). Plants raised from Zn-primed seeds and inoculated with *Azospirillum* had the higher P in grain (Fig. 6). When plants from Zn-primed seeds inoculated with *Azospirillum* grain P content was increased up to 6.8%.

Discussion

Soaking seeds in water prior to sowing followed by surface-drying (on-farm seed priming) can be simultaneously combined with micro-dosing of fertilizer, small amount of micronutrients, plant growth regulators and water restricting agents (polyethylene glycol). In this study, simultaneously to priming, wheat seeds were treated with Zn, Mn or IAA added to the conditioning solution. Seeds primed with water, auxin, Mn and, especially Zn solution resulted in increased grain weight, spike no per plant and seed no per spike. Plants of the Zn-primed seeds among different seed priming agents had the highest grain yield and yield components. Harris et al. [30], [53, 54], [23] and [10] reported that on-farm, soaking seeds in water prior to sowing resulted in faster emergence, higher vigor and density, accelerated flowering and maturity and higher yields as compared with non-prime crops. Considerable studies have shown that priming seeds with micronutrients can improve yield of crops [23]. Working with wheat and




Fig. 6 The effect of priming seeds for 6 h in water, IAA (2 ppm), or with aqueous solutions of Zn (0.2%) or Mn (0.2%) in combination or not with *Azospirillum* inoculation on grain N and P contents. Values

with the same letters are not significantly different at $P < 0.05$. Values are mean \pm standard error (SE) ($n = 3$)

barley crops under field conditions, [33] demonstrated that grain yield increased by on-farm seed priming with zinc. Likewise, Ajouri et al. [1] found that priming barley seeds with Zn enhanced seed germination and seedling development. Harris et al. [31] reported that seed priming with ZnSO₄ increased grain yield of wheat. At research and Extension Center, Arkansas, USA, [80] treated rice (*Oryza sativa* L.) seeds with Zn and found that growth and grain yield were increased. Micronutrients such as zinc were evaluated as cost-effective in several crops and suggested to be a promising alternative to soil- or foliar-applied Zn. Johnson et al. [37] found that grain yield of chickpea, lentil, rice and wheat decreased or was not affected, when seeds were primed with Zn.

In the present study, Mn priming treatment had also potential to improve grain yield of wheat, but at the rate less than that achieved by Zn-seed priming. In Australia, [49] soaked wheat seeds in aqueous solution of MnSO₄ and reported that grain yield and grain Mn content were increased. Priming seeds with MnSO₄ has been proposed to be the effective in improving wheat yield [23].

Since in this study seeds were primed with IAA, therefore improved yield components and grain yield can be related to the effect of auxin in the early growth of seedlings. In the present study, dry weights of plants of auxin-primed seeds were increased over non-primed seedling plants. It seems that auxin absorbed by seeds through priming could modulate plant processes. Auxin induces the root formation and coleoptile elongation (Taiz and Zeiger [84]). Diverse plant processes such as cell elongation and root initiation and development [60], [71] are regulated by auxin. Even regeneration of xylem and development of vascular development are modulated by auxin (Mattsson [48]). Partitioning of photo-assimilate from vegetative organs into reproductive organ is affected

by the auxin (Cole and Patrick [15]), an important factor causing grain yield to increase.

In the present study, neither field emergence nor emergence rate was significantly affected by seed priming treatments for both years of the study (data not shown) but seedlings fresh weights from primed seeds with different priming agents (Zn, Mn, IAA and water) were significantly higher than non-primed seeds after 30 and 60 days of emergence. In this experiment, treatments that improved seedling growth during early plant growth also increased yield components (spike no, 1000-grains weight and harvest index) and total grain yield, suggesting that increment in yield due to seed priming can be attributed to the improved seedling growth. Ashraf and Foolad [2] and Chen and Arora [13] indicated that changes in important metabolic pathways, exerting during early plant growth by seed priming, are able to continue until later stages of plant development. Semiarid area is characterized by irregular rainfall pattern, low and unpredictable rain conditions. This study was performed in a semiarid area, with irregular and insufficient rainfall during both years of the experiments. Virk et al. [90] and [59] attributed greater yield increments to the rainfall profiles, rain seasons with restricted precipitation resulted in greater yield increments. Additionally, early growth and development of seedlings from primed seeds allow plant to access and take up more fertilizer [30]. Any delay in development of seedling leads the fertilizer to be leached or volatilized from soil surface.

In this study, *Azospirillum*-inoculated plants produced more grain yield than non-inoculated seeds. Plants inoculated with *Azospirillum* produced grains with higher amount of N and P content. When inoculation was coupled with priming, additional increase in grain yield and N and P content in grains was exhibited. The benefits of inoculation on enhanced growth and yield could be attributed to

the phytohormone production, N₂ fixation, and P solubilization. Several literature regarding the promoting effects of *Azospirillum* on yield and other agronomic traits in wheat has been published previously [74], [65, 66], [4], [35], [38]. Increased phosphorus contents in grains due to *Azospirillum* inoculation could not merely refer to inoculation since the used bacterial strain in this study exhibited low potential activity of solubilizing inorganic phosphate in the modified Pikovskaya medium (data not shown). Higher N content in grains followed by inoculation could be attributed to some extent to the biological nitrogen fixation activity of this strain. In this study, inoculation resulted in attaining the early growth promotion in wheat seedlings. Auxin produced by *Azospirillum* has been assumed to have a promoting effect on attaining the early growth promotion in wheat [40]. Auxin induces rooting, root initiation, root development [60], [71] and increases root surface area [34]. Developed root system assists the plant in enhancing nutrient absorption [20], Gray and Smith [27], [83], [5] which can be important under dryland farming [38], Zarea [97]. Developed root system assists plant in uptaking soil minerals and water in a better way [58], [64]. In the present study, additional increase in growth and yield occurred when seed priming was combined with *Azospirillum*. Relationships of environment, *Azospirillum* and environment are complex. Various agronomic management methods such as applying N fertilizers and priming seed with chemical agent like 2,4-D and hydrogen peroxide have been reported to improve promoting effect of *Azospirillum* on plants [96].

Conclusions

It should be noted that there are a few reports dealing with PGPR inoculation coupled with priming in the field study. PGPR can be applied simultaneously to priming or following by priming. In the present study, *Azospirillum* inoculation following by seed priming is a significant additional approach in terms of grain yield and grain nutrient content.

Compliance with Ethical Standards

Conflict of interest The authors declare that they have no conflict of interest.

References

1. Ajouri A, Asgedom H, Becker M (2004) Seed priming enhances germination and seedling growth of barley under conditions of P and Zn deficiency. *J Plant Nutr Soil Sci* 167:630–636
2. Ashraf M, Foolad MR (2005) Pre-sowing seed treatment—a shotgun approach to improve germination, plant growth, and crop yield under saline and non-saline conditions. *Adv Agron* 88:223–271
3. Aymen EM, Kaouther Z, Fredj MB, Cherif H (2012) Seed priming for better growth and yield of safflower (*Carthamus tinctorius*) under saline condition. *J Stress Physiol Biochem* 8:135–143
4. Ayyaz K, Zaheer A, Rasul G, Mirza MS (2016) Isolation and identification by 16S rRNA sequence analysis of plant growth-promoting *Azospirillum* from the rhizosphere of wheat. *Braz J Microbiol* 47:542–550
5. Bashan Y, Holguin G, de Bashan LE (2004) *Azospirillum*–plant relationship: physiological, molecular, agriculture, and environmental advances (1997–2003). *Can J Microbiol* 50:521–577
6. Beijerinck MW (1925) Über ein *Spirillum*, welches freien Stickstoff binden kann. *Zentralbl Bakteriol Parasitenkd Infekt Abt.* 63:353
7. Bradford M (1976) A rapid and sensitive method for the quantitation of microgram quantities of protein utilizing the principle of protein-dye binding. *Anal Biochem* 72:248–254
8. Bradford KJ (1986) Manipulation of seed water relations via osmotic priming to improve germination under stress conditions. *HortSci*. 21:1105–1112
9. Cakmak I (2008) Enrichment of cereal grains with zinc: agronomic or genetic biofortification? *Plant Soil* 302:1–17
10. Carrillo-Reche J, Vallejo-Marín M, Quilliam RS (2018) Quantifying the potential of ‘on-farm’ seed priming to increase crop performance in developing countries. A meta-analysis. *Agron Sustain Dev* 38:64
11. Chapman HD, Pratt PF (1961) Ammonium vanadate–molybdate method for determination of phosphorus. In: Chapman HD, Pratt PF (eds) *Methods of analysis for soils. Plants and Water* University of California, Berkeley, pp 184–203
12. Chasapis CT, Loutsidou AC, Spiliopoulou CA, Stefanidou ME (2012) Zinc and human health: an update. *Arch Toxicol* 86:521–553
13. Chen K, Arora R (2013) Priming memory invokes seed stress-tolerance. *Environ Exp Bot* 94:33–45
14. Chen P, Bornhorst J, Aschner M (2018) Manganese metabolism in humans. *Front Biosci (Landmark Ed)*. 23:1655–1679
15. Cole DMA, Patrick JW (1998) Auxin control of photoassimilate transport to and within developing grains of wheat. *Aust J Plant Physiol* 25:69–77
16. Copur O, Demirel U, Karakus M (2010) Effects of several plant growth regulators on the yield and fiber quality of cotton (*Gossypium hirsutum* L.). *Notulae Botanicae HortiAgrobotanici Cluj* 38:104–110
17. Curtin D, Martin RJ, Scott CL (2008) Wheat (*Triticum aestivum*) response to micronutrients (Mn, Cu, Zn, B) in Canterbury, New Zealand. *New Zealand J Crop Hortic Sci* 36:169–181
18. de Bashan LE, Mayali X, Bebout BM, Weber PK, Detweiler AM, Hernandez JP, Prufert-Bebout L, Bashan Y (2016) Establishment of stable synthetic mutualism without co-evolution between microalgae and bacteria demonstrated by mutual transfer of metabolites (NanoSIMS isotopic imaging) and persistent physical association (fluorescent in situ hybridization). *Algal Res* 15:179–186
19. de Souza R, Ambrosini A, Passaglia LM (2015) Plant growth-promoting bacteria as inoculants in agricultural soils. *Genet Mol Biol.* 38:401–419
20. Dey R, Pal KK, Bhatt DM, Chauhan SM (2004) Growth promotion and yield enhancement of peanut (*Arachis hypogaea* L.) by application of plant growth promoting rhizobacteria. *Microbiol Res* 159:371–394
21. Dotto L, Silva VN (2017) Beet seed priming with growth regulators. *Semina: CienciasAgrarias* 38:1785–1798

22. Farahmandfar E, Shirvan MB, Sooran SA, Hoseinzadeh D (2013) Effect of seed priming on morphological and physiological parameters of fenugreek seedlings under salt stress. *Intl J Agri Crop Sci* 5:811–815
23. Farooq M, Basra SMA, Wahid A (2006) Priming of field-sown rice seed enhances germination, seedling establishment, allometry and yield. *Plant Growth Regul* 49:285–294
24. Gao Y, Wu P, Zhao X, Wang Z (2014) Growth, yield, and nitrogen use in the wheat/maize intercropping system. *Field Crops Res* 167:19–30
25. Garvin DF, Welch RM, Finley JW (2006) Historical shifts in the seed mineral micronutrient concentration of US hard red winter wheat germplasm. *J Sci Food Agric* 86:2213–2220
26. Gibson RS (2006) Zinc: the missing link in combating micronutrient malnutrition in developing countries. *Proc Nutr Soc* 65:51–60
27. Gray EJ, Smith DL (2005) Intracellular and extracellular PGPR: Commonalities and distinctions in the plant-bacterium signaling process. *Soil Biol Biochem* 37:395–412
28. Gulluoglu L (2004) Determination of usage of plant growth regulators in soybean (*Glycine max* Merr) farming under Harran plain conditions. *J Fac. Agri* 8:17–23
29. Hafeez B, Khanif YM, Saleem M (2013) Role of zinc in plant nutrition—a review. *Am J Exp Agric* 3:374–391
30. Harris D, Raghuvanshi BS, Gangwar JS, Singh SC, Joshi KD, Rashid A, Hollington PA (2001) Participatory evaluation by farmers of “on-farm” seed priming in wheat in India. *Exp Agric* 37:403–415
31. Harris D, Rashid A, Arif M, Yunas M (2005) Alleviating micronutrient deficiencies in alkaline soils of the North-West Frontier Province of Pakistan: Onfarm seed priming with zinc in wheat and chickpea. In: Andersen P, Tuladhar JK, Karki KB, Maskey SL (eds) *Micronutrients in South and South East Asia*, pp 143–151
32. Harris D, Rashid A, Miraj G, Arif M, Shah H (2007) On-farm seed priming with zinc sulphate solution—a cost-effective way to increase the maize yields of resource-poor farmers. *Field Crops Res*. 10:119–127
33. Harris D, Rashid A, Miraj G, Arif M, Yunas M (2008) ‘On-farm’ seed priming with zinc in chickpea and wheat in Pakistan. *Plant Soil* 306:3–10
34. Ivanchenko MG, Napsucially-Mendivil S, Dubrovsky JG (2010) Auxin-induced inhibition of lateral root initiation contributes to root system shaping in *Arabidopsis thaliana*. *Plant J* 64:740–752
35. Jafarian T, Zarea MJ (2016) Hydrogen peroxide affects plant growth promoting effects of *Azospirillum*. *Crop Sci Biotech* 19:167–175
36. Jafri N, Mazid M, Mohammad F (2015) Responses of seed priming with gibberellic acid on yield and oil quality of sunflower (*Helianthus annuus* L.). *Indian J. Agric. Res.* 49:235
37. Johnson SE, Lauren JG, Welch RM, Duxbury JM (2005) A comparison of the effects of micronutrient seed priming and soil fertilization on the mineral nutrition of chickpea (*Cicer arietinum*), lentil (*Lens culinaris*), rice (*Oryza sativa*) and wheat (*Triticum aestivum*) in Nepal. *Exp Agric* 41:427–448
38. Karimi N, Zarea MJ, Mehnaz S (2018) Endophytic *Azospirillum* for enhancement of growth and yield of wheat. *Environ Sustain* 1:149–158
39. Khalid BY, Malik NSA (1982) Pre-sowing soaking of wheat seeds in copper and manganese solutions. *Commun Soil Sci Plant Analy* 13:981–986
40. Khalid A, Arshad M, Zahir ZA (2004) Screening plant growth-promoting rhizobacteria for improving growth and yield of wheat. *J Appl Microbiol* 96:473–480
41. Khammas KM, Ageron E, Grimont PAD, Kaiser P (1989) *Azospirillum irakense* sp. nov., a nitrogen-fixing bacterium associated with rice roots and rhizosphere soil. *Res Microbiol* 140:679–693
42. Kopylov IP, Spirydonov VH, Patyka VP (2009) Identification of *Azospirillum* genus bacteria isolated from the spring wheat root zone. *Mikrobiol Z* 71:13–19
43. Laity JH, Lee BM, Wright PE (2001) Zinc finger proteins: new insights into structural and functional diversity. *Curr Opin Struct Biol* 11:39–46
44. Lecube ML, Noriega GO, Santa Cruz DM, Tomaro ML, Batlle A, Balestrasse KB (2014) Indole acetic acid is responsible for protection against oxidative stress caused by drought in soybean plants: the role of heme oxygenase induction. *Redox Rep* 19:242–250
45. Liu DH, Jiang WS, Wuang C (1996) Effects of Zn²⁺ on root growth, cell division, and nucleoli of *Allium cepa* L. *J Environ Sci* 8:21–27
46. Ma T, Duan XH, Yang YY, Yao J, Gao TP (2017) Zinc-alleviating effects on iron-induced phytotoxicity in roots of *Triticum aestivum*. *Biol Plant* 61:733–740
47. Ma HY, Zhao DD, Ning QR, Wei JP, Li Y, Wang MM, Liu XL, Jiang Ch-J (2018) A multi-year beneficial effect of seed priming with gibberellic acid-3 (GA₃) on plant growth and production in a perennial grass. *Leymus Chinensis Sci Rep* 8:13214
48. Mattsson J, Ckurshumova W, Berleth T (2003) Auxin signaling in Arabidopsis leaf vascular development. *Plant Physiol* 131:1327–1339
49. Marcar NE, Graham RD (1986) Effect of seed manganese content on the growth of wheat (*Triticum aestivum*) under manganese deficiency. *Plant Soil* 96:165–173
50. Mehnaz S, Mirza MS, Haurat J, Bally R, Normand P, Bano A, Malik KA (2001) Isolation and 16S rRNA sequence analysis of the beneficial bacteria from the rhizosphere of rice. *Can J Microbiol* 47:110–117
51. Mehnaz S, Weselowski B, Lazarovits G (2007) *Azospirillum zeae* sp. nov., diazotrophic bacteria isolated from rhizosphere soil of *Zea mays*. *Int J Syst Evol Microbiol* 57:2805–2809
52. Mehnaz S, Weselowski B, Lazarovits G (2007) *Azospirillum canadense* nov., a nitrogen fixing bacterium isolated from corn rhizosphere. *Int J Syst Evol Microbiol* 57:620–624
53. Murungu FS, Nyamugafata P, Chiduzo C et al (2003) Effects of seed priming, aggregate size and soil matric potential on emergence of cotton (*Gossypium hirsutum* L.) and maize (*Zea mays* L.). *Soil Tillage Res* 74:161–168
54. Murungu FS, Chiduzo C, Nyamugafata P, Clark LJ, Whalley WR (2004) Effect of on-farm seed priming on emergence, growth and yield of cotton and maize in a semi-arid area of Zimbabwe. *Exp Agric* 40:23
55. Naeem M, Bhatti I, Ahmad RH, Ashraf MY (2004) Effect of some growth hormones (GA₃, IAA and Kinetin) on the morphology and early or delayed initiation of bud of lentil (*Lens culinaris* Medik). *Pak J Bot* 36:801–809
56. Naidoo G, Chirkoot HD (2004) The effects of coal dust on photosynthetic performance of the mangrove, *Avicennia marina* in Richards Bay, South Africa. *Environ Pollut* 127:359–366
57. Okon Y, Kapulnik Y (1986) Development and function of *Azospirillum*-inoculated roots. *Plant Soil* 90:3–16
58. Okon Y, Bloembergen GV, Lugtenberg BJJ (1998) Biotechnology of biofertilization and phytostimulation. In: Altman A (ed) *Agricultural biotechnology*. Marcel Dekker, New York, pp 327–349
59. Ousman A, Aune JB (2011) Effect of seed priming and micro-dosing of fertilizer on groundnut, sesame and cowpea in Western Sudan. *Exp Agric* 47(3):431–443
60. Overvoorde P, Fukaki H, Beeckman T (2010) Auxin control of root development. *Cold Spring Harb Perspect Biol* 2(6):a001537

61. Padole VR (1979) Effect of presoaking seed treatment of wheat seed with chemicals and hormones on yield and uptake of NPK. *J Maharashtra AgricUniv* 4(1):85–88
62. Pallaoro DS et al (2016) Priming corn seeds with plant growth regulator. *J Seed Sci* 38:227–232
63. Palmer CM, Guerinot ML (2009) Facing the challenges of Cu, Fe and Zn homeostasis in plants. *Nat Chem Biol* 5:333–340
64. Pérez-Montaña F, Alías-Villegas C, Bellogín RA, del Cerro P, Espuny MR, Jiménez-Guerrero I, López-Baena FJ, Ollero FJ, Cubo T (2014) Plant growth promotion in cereal and leguminous agricultural important plants: from microorganism capacities to crop production. *Microbiol Res* 169:325–336
65. Piccinin GG, Dan LGM, Braccini AL, Mariano DC, Okumura RS, Bazo G, Ricci TT (2011) Agronomic efficiency of *Azospirillum brasilense* in physiological parameters and yield components in wheat crop. *J Agron* 10:132–135
66. Piccinin GG, Braccini AL, Dan LGM, Scapim CA, Ricci TT, Bazo GL (2013) Efficiency of seed inoculation with *Azospirillum brasilense* on agronomic characteristics and yield of wheat. *Ind Crop Prod* 43:393–397
67. Rashid A (1996) Secondary and micronutrients. In: Bashir E, Bantel R (eds) *Soil science*. National Book Foundation, Islamabad, Pakistan, p 374
68. Reis S, Pavia I, Carvalho A, Moutinho-Pereira J, Correia C, Lima-Brito J (2018) Seed priming with iron and zinc in bread wheat: effects in germination, mitosis and grain yield. *Protoplasma* 255:1179–1194
69. Rengel Z, Graham RD (1995) Importance of seed Zn content for wheat grown on Zn-deficient soil I vegetative growth. *Plant Soil* 173:259–266
70. Rengel Z, Graham RD (1995) Importance of seed Zn content for wheat grown on Zn-deficient soil II grain yield. *Plant Soil* 173:267–274
71. Saini S, Sharma I, Kaur N, Pati PK (2013) Auxin: a master regulator in plant root development. *Plant Cell Rep* 32:741–757
72. Salantur A, Ozturk R, Akten S (2006) Growth and yield response of spring wheat (*Triticum aestivum* L.) to inoculation with rhizobacteria. *Plant Soil Environ*. 52:111–118
73. Salisbury FB (1994) The role of plant hormones. In: Wilkinson RE (ed) *Plant environment interactions*. Dekker, New York
74. Santa ORD, Hernadez RF, Alvarez GLM, Ronzelli P, Soccol CR (2004) *Azospirillum* sp. inoculation in wheat, barley and oats seeds greenhouse experiments. *Braz Arch Biol Technol* 47:843–850
75. Saraswat VK, Bansal KN (1991) Methods of zinc application and its effect on yield and zinc content of rice (*Oryza sativa*) and wheat (*Triticum vulgare*). *Madras Agric J* 78(5–8):174–177
76. Schmidt SB, Jensen PE, Husted S (2016) Manganese deficiency in plants: the impact on photosystem II. *Trends Plant Sci* 21:622–632
77. Shetiwy M, Shen H, Xu J, Guan Y, Song W, Hu J (2017) Seed polyamines metabolism induced by seed priming with spermidine and 5-amino levulinic acid for chilling tolerance improvement in rice (*Oryza sativa* L.) seedlings. *Environ Exp Bot* 137:58–72
78. Singh B, Natesan SKA, Singh BK, Usha K (2003) Improving zinc efficiency of cereals under zinc deficiency. *Curr Sci* 88:36–44
79. Singh SP, Keller B, Gruissem W, Bhullar NK (2017) Rice nicotianamine synthase 2 expression improves dietary iron and zinc levels in wheat. *Theor Appl Genet* 130:283–292
80. Slaton NA, Wilson-Jr CE, Ntamatungiro S, Norman RJ, Boothe DL (2001) Evaluation of zinc seed treatments for rice. *Agron J* 93:152–157
81. Steenhoudt O, Vanderleyden J (2000) *Azospirillum*, a free-living nitrogen-fixing bacterium closely associated with grasses: genetic, biochemical and ecological aspects. *FEMS Microbiol* 24:487–506
82. Stout PR, Arnon DI (1939) Experimental methods for the study of the role of copper, manganese, and zinc in the nutrition of higher plants. *Am J Bot* 26:144–149
83. Sukumar P, Legué V, Vayssières A, Martin F, Tuskan GA, Kalluri UC (2013) Involvement of auxin pathways in modulating root architecture during beneficial plant–microorganism interactions. *Plant, Cell Environ* 36:909–919
84. Taiz L, Zeiger E (2010) *Plant Physiology*, 5th edn. Sinauer Associates, Inc., Publishers, Sunderland, MA, USA
85. Tarrand JJ, Krieg NR, Döbereiner J (1978) A taxonomic study of the *Spirillum lipoferum* group, with the descriptions of a new genus, *Azospirillum* gen. nov. and two species *Azospirillum lipoferum* (Beijerinck) comb. nov. and *Azospirillum brasilense* sp. nov. *Can J Microbiol* 24:967–980
86. Teale WD, Paponov IA, Palme K (2006) Auxin in action: signalling, transport and the control of plant growth and development. *Nat Rev Mol Cell Biol* 7:847–859
87. Ulfat A, Majid SA, Hameed A (2017) Hormonal seed priming improves wheat (*Triticum aestivum* L.) field performance under drought and non-stress conditions. *Pak J Bot* 49:1239–1253
88. Vallee BL, Falchuk KH (1993) The biochemical basis of zinc physiology. *Physiol Rev* 73:79–118
89. Van Schouwenberg J, Walinge I (1973) *Methods of analysis for plant material*. Agricultural University, Wageningen
90. Virk DS, Chakraborty M, Ghosh J, Harris D (2006) Participatory evaluation of horsegram (*Macrotyloma uniflorum*) varieties and their on-station responses to on-farm seed priming in eastern India. *Exp Agric* 42:411
91. Wei LX et al (2017) Priming of rice (*Oryza sativa* L.) seedlings with abscisic acid enhances seedling survival, plant growth, and grain yield in saline-alkaline paddy fields. *Field Crops Res*. 203:86–93
92. White JG, Zasoski RJ (1999) Mapping soil micronutrients. *Field Crops Res* 60:11–26
93. Yilmaz A, Ekiz H, Gultekin I, Torun B, Barut H, Karanlik S, Cakmak I (1998) Effect of seed zinc content on grain yield and zinc concentration of wheat grown in zinc-deficient calcareous soils. *J Plant Nutr* 21:2257–2264
94. Zahid M, Abbasi MK, Hameed S, Rahim N (2015) Isolation and identification of indigenous plant growth promoting rhizobacteria from Himalayan region of Kashmir and their effect on improving growth and nutrient contents of maize (*Zea mays* L.). *Front Microbiol* 6:207
95. Zahid M, Abbasi MK, Hameed S, Rahim N (2015) Isolation and identification of indigenous plant growth promoting rhizobacteria from Himalayan region of Kashmir and their effect on improving growth and nutrient contents of maize. *Front Microbiol* 6:207
96. Zarea MJ (2017) *Azospirillum* and wheat production. In: Kumar V, Kumar M, Sharma SH (eds) *Probiotics in agroecosystem*. Springer, Singapore, pp 329–348
97. Zarea MJ (2019) Applications of Beneficial Microbe in Arid and Semiarid Agroecosystem: IAA-Producing Bacteria. In: Kumar V, Prasad R, Kumar M, Choudhary D (eds) *Microbiome in Plant Health and Disease*. Springer, Singapore, pp 105–118
98. Zarea MJ (2019) Salt-tolerant microbes: isolation and adaptation. In: Varma A, Giri B (eds) *Microorganisms in saline environments strategies and functions*. Springer, Switzerland, pp 285–301

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.